


## geometry of diagonalization



$$\vec{b}_1 = \langle 1, 1 \rangle, \vec{b}_2 = \langle -2, 1 \rangle \quad B = \begin{bmatrix} 1 & -2 \\ 1 & 1 \end{bmatrix} \quad B^{-1} = \frac{1}{3} \begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix}$$

$$\vec{x} = B \vec{y} : \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 1 & -2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} \quad \begin{aligned} x_1 &= y_1 - 2y_2 \\ x_2 &= y_1 + y_2 \end{aligned}$$

$$\vec{y} = B^{-1} \vec{x} : \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \quad \begin{aligned} y_1 &= \frac{1}{3}(x_1 + 2x_2) \\ y_2 &= \frac{1}{3}(-x_1 + x_2) \end{aligned}$$


$$A = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$$

→ (right click menu) →

$$A_B = B^{-1} A B = \frac{1}{3} \begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 1 & 1 \end{bmatrix}$$

$$= \frac{1}{3} \begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 5 & 2 \\ 5 & -1 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 15 & 0 \\ 0 & -3 \end{bmatrix} = \begin{bmatrix} 5 & 0 \\ 0 & -1 \end{bmatrix} \checkmark$$

Linear Algebra: Eigenvectors (A) =  $\begin{bmatrix} -1 \\ 5 \end{bmatrix}, \begin{bmatrix} -2 & 1 \\ 1 & 1 \end{bmatrix}$  ordering of eigenvalues still random



columns are corresponding eigenvectors in same order

$$\lambda_1 = 5, \vec{b}_1 = \langle 1, 1 \rangle \\ \lambda_2 = -1, \vec{b}_2 = \langle -2, 1 \rangle$$

# 1st order linear homogeneous DE system : real eigenvalues

$$A = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} \quad \lambda = 5, -1 \quad B = \begin{bmatrix} 1 & -2 \\ 1 & 1 \end{bmatrix} \quad B^{-1} = \frac{1}{3} \begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix} \quad A_B = B^{-1}AB = \begin{bmatrix} 5 & 0 \\ 0 & -1 \end{bmatrix}$$

$$\vec{x}' = A\vec{x} \quad \frac{dx_1}{dt} = x_1 + 4x_2 \quad \frac{dx_2}{dt} = 2x_1 + 3x_2 \quad \vec{x} = B\vec{y} \quad \vec{y}' = A_B \vec{y} \quad \frac{dy_1}{dt} = 5y_1 \rightarrow y_1 = c_1 e^{5t}$$

$$\vec{y} = B^{-1}\vec{x} \quad \frac{dy_2}{dt} = -y_2 \rightarrow y_2 = c_2 e^{-t}$$

$\downarrow$

$$\vec{x}(0) = \begin{bmatrix} -2 \\ 4 \end{bmatrix} \rightarrow x_1(0) = -2, x_2(0) = 4 \quad y(0) = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \leftarrow \quad y_1(0) = c_1, y_2(0) = c_2$$

$$\vec{x}(0) = B\vec{y}(0) \rightarrow \begin{bmatrix} -2 \\ 4 \end{bmatrix} = B \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \rightarrow \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = B^{-1} \begin{bmatrix} -2 \\ 4 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$

general soln:

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 1 & -2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} c_1 e^{5t} \\ c_2 e^{-t} \end{bmatrix} = \begin{bmatrix} c_1 e^{5t} - 2c_2 e^{-t} \\ c_1 e^{5t} + c_2 e^{-t} \end{bmatrix}$$

$$= c_1 e^{5t} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 e^{-t} \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

↑ growth along  $\vec{b}_1$       ↑ decay along  $\vec{b}_2$

IVP Soln:


$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2e^{5t} - 4e^{-t} \\ 2e^{5t} + 2e^{-t} \end{bmatrix}$$

$$= 2e^{5t} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 2e^{-t} \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

eigenvector entries set relative initial values for variables in each "mode" of exponential behavior

initial data sets the "mixture" coefficients of the combination of modes

the eigenvalues are the exponential rate coefficients for each mode



the decaying mode ( $\lambda = -1$ ) decays to 1% of its original value in the time interval  $t = 0..4.6$  during which the growing mode ( $\lambda = 5$ ) grows by a factor  $e^{5(4.6)} \approx 10^{10}$  (very big)