

# Calculus: Basic Functions for instant recall

	diff	int	simple u-sub
power	$\frac{d}{dx} x^n = n x^{n-1}$	$\int x^n dx = \frac{x^{n+1}}{n+1} + C$	$\int (ax+b)^n dx = \frac{(ax+b)^{n+1}}{a(n+1)} + C$
ln	$\frac{d}{dx} \ln x  = \frac{1}{x}$	$\int \frac{1}{x} dx = \ln x  + C$	$\int \frac{1}{ax+b} dx = \frac{1}{a} \ln ax+b  + C$
exp	$\frac{d}{dx} e^x = e^x$	$\int e^x dx = e^x + C$	$\int e^{ax} dx = \frac{1}{a} e^{ax} + C$
trig	$\frac{d}{dx} \sin x = \cos x$ $\frac{d}{dx} \cos x = -\sin x$	$\int \cos x dx = \sin x + C$ $\int \sin x dx = -\cos x + C$	$\int \cos ax dx = \frac{1}{a} \sin ax + C$ $\int \sin ax dx = -\frac{1}{a} \cos ax + C$
chain/ u-sub	$\frac{d}{dx} f(u) = \underbrace{f'(u)}_{\frac{df}{du}} \frac{du}{dx}$	$\int f(u(x)) \frac{du(x)}{dx} dx = \int f(u) du = F(u) + C = F(u(x)) + C$ if $F'(x) = f(x)$ "antiderivative"	
additive constant	$\frac{d}{dx} (f(x) + c) = \frac{d}{dx} f(x)$	$\int c f(x) dx = c \int f(x) dx$	} + ≠ * !
multiplicative constant	$\frac{d}{dx} (c f(x)) = c \frac{d}{dx} f(x)$		

You are expected to be able to do any of the above explicit integrals by hand, or any that can be reduced to them by an obvious u-substitution. Any derivative or integral you are uncertain of you are expected to check symbolically with Maple or your graphing calculator. There is no excuse for getting a derivative or integral wrong with technology at your fingertips.

MAT 2500 = CALC 3 and MAT 2705 = DE w/ Lin Alg assume these basic operations from CALC 1 and CALC 2 and build on them.

## Some Algebra Rules

distributive rule:  $a(b+c) \xrightarrow{\text{expand}} ab+ac$  [  $a+b = a \cdot 1 + b = 1(a+b)$  ]  
 $\xleftarrow{\text{factor}}$

fractions:

$$\frac{a}{b} + \frac{c}{d} = \frac{ad+bc}{bd} = \frac{ad+bc}{bd}$$

$$\frac{a}{b} + c = \frac{a}{b} + \frac{bc}{b} = \frac{a+bc}{b}$$

$$a^{-1} = \frac{1}{a}, \left(\frac{a}{b}\right)^{-1} = \frac{b}{a}$$

$$\frac{c}{\left(\frac{a}{b}\right)} = \left(\frac{b}{a}\right)c = \frac{bc}{a}$$

add numerators  
if same  
denominator

(cannot cancel  
b's in last expression)

$$\left(\frac{a}{b}\right) \frac{c}{c} = \frac{a}{bc}$$

always factor before  
cancelling:  
 $\frac{a+ac}{ab} = \frac{a(1+c)}{a \cancel{b}} = \frac{1+c}{b}$

cancellation:

$$*: \frac{ab^1}{b^1} = a$$

$$+: a + \cancel{b} = \cancel{b} + a$$

only base e  
needed:

$$a = e^{\ln a}$$

$$a^x = (e^{\ln a})^x = e^{(\ln a)x}$$

$$\log_a x = \frac{\ln x}{\ln a}$$

exponentials ( $a > 0$ ):

logs ( $\ln = \log_e$ ):

product  $a^x a^y = a^{x+y}$

$\ln xy = \ln x + \ln y$

quotient  $a^x / a^y = a^{x-y}$

$\ln x/y = \ln x - \ln y$

power  $(a^x)^y = a^{xy}$

$\ln x^p = p \ln x$

reciprocal  $(a^x)^{-1} = a^{-x} = 1/a^x$

$\ln \frac{1}{x} = \ln x^{-1} = -\ln x$

exp/ln properties:  $e^{\ln x} = x$  ( $x > 0$ )  $e^0 = 1$   $e^1 = e \approx 2.718$

$\ln e^x = x$   $\ln 1 = 0$

[  $e^{p \ln x} = (e^{\ln x})^p = x^p$  ]  $\frac{1}{e^x} = e^{-x}$  (preferred)

powers / roots:

product  $(xy)^p = x^p y^p$

$x^0 = 1$  ( $x \neq 0$ )

quotient  $(x/y)^p = x^p / y^p$

$1^x = 1$

power  $(x^p)^q = x^{pq}$

$\sqrt{x^2} = |x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$

reciprocal  $x^{-p} = 1/x^p$

$x^{\frac{1}{n}} = \sqrt[n]{x}$ ,  $n$  integer  $> 0$

always convert radical notation to power notation to work with in calculus

$x^{\frac{m}{n}} = (x^{\frac{1}{n}})^m = (\sqrt[n]{x})^m$

$= (x^m)^{\frac{1}{n}} = \sqrt[n]{x^m}$

$(a+b)^2 \xrightarrow{\text{expand}} a^2 + 2ab + b^2$   
 $\xleftarrow{\text{factor}}$

(binomial theorem for higher integer powers)

## Some Algebra Rules 2

solving equations

a) linear:  $ax+b=0 \rightarrow x = -b/a$  ( $a \neq 0$ )

b) quadratic:  $ax^2+bx+c=0 \rightarrow x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$  ( $a \neq 0$ ) memorize!

related technique: completing the square

$$ax^2+bx+c = a\left(x^2 + \frac{b}{a}x\right) + c$$

$$\left(x + \frac{b}{2a}\right)^2 - \left(\frac{b}{2a}\right)^2 \quad \left[\text{since } x^2 + 2x\left(\frac{b}{2a}\right) + \left(\frac{b}{2a}\right)^2 - \left(\frac{b}{2a}\right)^2\right]$$

$$= a\left(x + \frac{b}{2a}\right)^2 - a\left(\frac{b^2}{4a^2}\right) + c$$

$$= a\left(x + \frac{b}{2a}\right)^2 + \left(c - \frac{b^2}{4a}\right)$$

don't memorize formula  
remember technique

Distinguish between "solving an equation" for an unknown variable which appears in it, as opposed to "simplifying or rewriting" an expression, as in:

$$x^2 - 2x = x(x-2) \quad (\text{factored})$$

Note:  $f(x)g(x) = 0 \rightarrow$  either  $f(x) = 0$  or  $g(x) = 0$

$\frac{f(x)}{g(x)} = 0 \rightarrow f(x) = 0$ , provided that  $g(x) \neq 0$  at a solution of  $f(x) = 0$

rules of algebra NOT!

$\sqrt{a+b} \neq \sqrt{a} + \sqrt{b}$ ,  $\sqrt{a^2+b^2} \neq a+b$

cancellation NOT!

$\frac{c}{a+b} \neq \frac{c}{a} + \frac{c}{b}$ ,  $\frac{ab+c}{ad} \neq \frac{b+c}{d}$

\*:  $\frac{a+bc}{b} \neq a+c$

$\sqrt{x^2} \neq x$  unless  $x$  is known to be positive ( $|x|$ !)

+:  $b + a(c-b) \neq ac$

$f(2x) \neq 2f(x)$  (unless  $f$  is very special!)

$\sin 2x \neq 2 \sin x$

Miscellaneous

$0x = 0$  (says nothing about  $x$ )

even roots of negative #'s are complex, not real:

$\frac{0}{x} = 0$  when  $x \neq 0$

$\sqrt[n]{x}$ ,  $x < 0$  is undefined over real #'s for  $n > 0$  integer

$\frac{x}{0}$  never defined

maths case sensitive:  $T \neq t$ ,  $R \neq r$