

Calculus: Basic Functions for instant recall

	diff	int	simple u-sub
power	$\frac{d}{dx} x^n = n x^{n-1}$	$\int x^n dx = \frac{x^{n+1}}{n+1} + C$	$\int (ax+b)^n dx = \frac{(ax+b)^{n+1}}{a(n+1)} + C$
ln	$\frac{d}{dx} \ln x = \frac{1}{x}$	$\int \frac{1}{x} dx = \ln x + C$	$\int \frac{1}{ax+b} dx = \frac{1}{a} \ln ax+b + C$
exp	$\frac{d}{dx} e^x = e^x$	$\int e^x dx = e^x + C$	$\int e^{ax} dx = \frac{1}{a} e^{ax} + C$
trig	$\frac{d}{dx} \sin x = \cos x$ $\frac{d}{dx} \cos x = -\sin x$	$\int \cos x dx = \sin x + C$ $\int \sin x dx = -\cos x + C$	$\int \cos ax dx = \frac{1}{a} \sin ax + C$ $\int \sin ax dx = -\frac{1}{a} \cos ax + C$
chain/ u-sub	$\frac{d}{dx} f(u) = \underbrace{f'(u)}_{\frac{df}{du}} \frac{du}{dx}$	$\int f(u(x)) \frac{du(x)}{dx} dx = \int f(u) du = F(u) + C = F(u(x)) + C$ if $F'(x) = f(x)$ "antiderivative"	
additive constant	$\frac{d}{dx} (f(x) + c) = \frac{d}{dx} f(x)$	$\int c f(x) dx = c \int f(x) dx$	} + ≠ * !
multiplicative constant	$\frac{d}{dx} (c f(x)) = c \frac{d}{dx} f(x)$		

You are expected to be able to do any of the above explicit integrals by hand, or any that can be reduced to them by an obvious u-substitution. Any derivative or integral you are uncertain of you are expected to check symbolically with Maple or your graphing calculator. There is no excuse for getting a derivative or integral wrong with technology at your fingertips.

MAT 2500 = CALC 3 and MAT 2705 = DE w/ Lin Alg assume these basic operations from CALC 1 and CALC 2 and build on them.

Some Algebra Rules

distributive rule: $a(b+c) = ab + ac$

fractions: $\frac{a}{b} + \frac{c}{d} = \frac{ad+bc}{bd}$

$\frac{a}{b} + c = \frac{a+bc}{b}$

$a^{-1} = \frac{1}{a}$, $(\frac{a}{b})^{-1} = \frac{b}{a}$

$\frac{c}{(\frac{a}{b})} = \frac{bc}{a}$

expand
factor

$[a+ac = a(1+c)]$

add numerators
if same
denominator
(cannot cancel
b's in last expression)

always factor before
canceling:
 $\frac{a+ac}{ab} = \frac{a(1+c)}{ab} = \frac{1+c}{b}$

cancellation:
* $\frac{ab^1}{b^1} = a$
+ $a + b^{-1}b^0 = a$

only base e
needed:

exponentials ($a > 0$):

product $a^x a^y = a^{x+y}$

quotient $a^x / a^y = a^{x-y}$

power $(a^x)^y = a^{xy}$

reciprocal $(a^x)^{-1} = a^{-x} = 1/a^x$

logs ($\ln = \log_e$):

$\ln xy = \ln x + \ln y$

$\ln x/y = \ln x - \ln y$

$\ln x^p = p \ln x$

$\ln \frac{1}{x} = \ln x^{-1} = -\ln x$

$a = e^{\ln a}$

$a^x = (e^{\ln a})^x$

$= e^{(\ln a)x}$

$\log_a x = \frac{\ln x}{\ln a}$

exp/ln properties:

$e^{\ln x} = x$ ($x > 0$)

$\ln e^x = x$

$[e^{p \ln x} = (e^{\ln x})^p = x^p]$

$\frac{1}{e^x} = e^{-x}$ (preferred)

$e^1 = e \approx 2.718$

$\ln 1 = 0$

powers/roots:

product $(xy)^p = x^p y^p$

quotient $(x/y)^p = x^p / y^p$

power $(x^p)^q = x^{pq}$

reciprocal $x^{-p} = 1/x^p$

$x^0 = 1$ ($x \neq 0$)

$1^x = 1$

$\sqrt{x^2} = |x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$

$x^{\frac{1}{n}} = \sqrt[n]{x}$, n integer > 0

$x^{\frac{m}{n}} = (\sqrt[n]{x})^m = (\sqrt[n]{x^m})$

$(a+b)^2 = a^2 + 2ab + b^2$

(binomial theorem for higher integer powers)

Some Algebra Rules 2

solving equations

a) linear: $ax+b=0 \rightarrow x = -b/a$ ($a \neq 0$)

b) quadratic: $ax^2+bx+c=0 \rightarrow x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$ ($a \neq 0$) **MEMORIZE!**

related technique: completing the square

$ax^2+bx+c = a(x^2 + \frac{b}{a}x) + c$

$(x + \frac{b}{2a})^2 - (\frac{b}{2a})^2 + c$ [since $= x^2 + 2x(\frac{b}{2a}) + (\frac{b}{2a})^2$]

$= a(x + \frac{b}{2a})^2 - d(\frac{b^2}{4a^2}) + c$

$= a(x + \frac{b}{2a})^2 + (c - \frac{b^2}{4a})$

don't memorize formula
remember technique

Distinguish between "solving an equation" for an unknown variable which appears in it, as opposed to "simplifying or rewriting" an expression, as in:

$x^2-2x = x(x-2)$ (factored)

Note: $f(x)g(x)=0 \rightarrow$ either $f(x)=0$ or $g(x)=0$

$\frac{f(x)}{g(x)}=0 \rightarrow f(x)=0$, provided that $g(x) \neq 0$ at a solution of $f(x)=0$

Rules of algebra NOT!

$\sqrt{a+b} \neq \sqrt{a} + \sqrt{b}$, cancellation NOT!

$\frac{a}{b} + \frac{c}{d} \neq \frac{a+c}{b+d} \neq \frac{b+c}{a+d}$

$\sqrt{x^2} \neq x$ unless x is known to be positive ($|x|$)

$f(2x) \neq 2f(x)$ (unless f is very special!)

$\sin 2x \neq 2 \sin x$

cancellation NOT!

$x: \frac{a+bc}{b} \neq a+c$

$+: \frac{b+a(c-b)}{b} \neq ac$

Miscellaneous

$0 \times = 0$ (says nothing about x)

$\frac{0}{x} = 0$ when $x \neq 0$

$\frac{0}{0}$ never defined

maths case sensitive: $T \neq t, R \neq r$

even roots of negative #s are complex,
not real:

$\sqrt[n]{x}, x < 0$ is undefined over
real #s for $n > 0$ integer