


Show all work, including mental steps, in a clearly organized way that speaks for itself. Use proper mathematical notation, identifying expressions by their proper symbols (introducing them if necessary), and use arrows and equal signs when appropriate. Always simplify expressions. **BOX** final short answers. **LABEL** parts of problem. Keep answers EXACT (but give decimal approximations for interpretation). Indicate where technology is used and what type (Maple, GC, MathCad). You are encouraged to use technology to check all of your hand results.



The parametrized curve  $\vec{r}(t) = \langle t, t^2, 1 - t^2 \rangle$ ,  $-1 \leq t \leq 1$  is shown in the figure together with the vectors  $\vec{r}(-1)$ ,  $\vec{r}(0)$ ,  $\vec{r}(1)$ .

- a) Evaluate and simplify  $\vec{r}'(t)$ ,  $\vec{r}''(t)$ ,  $|\vec{r}'(t)|$ ,  $\vec{T}(t)$ .
- b) Evaluate  $\vec{r}(1)$ ,  $\vec{r}'(1)$ ,  $\vec{r}''(1)$ ,  $\vec{r}'(1) \times \vec{r}''(1)$ ,  $\vec{T}(1)$ .
- c) This is a plane curve which coincides with the plane containing the first and second derivatives. Write an equation for this plane picking any point on the curve and using a normal to the plane of the first and second derivatives at that point: simplify to a linear condition among the three coordinates.
- d) Write the parametrized equations of the line containing the points on the curve for which  $t=0$  and  $t=1$ , i.e.,  $x=...$ ,  $y=...$ ,  $z=...$ .
- e) What is the length of the straight line segment connecting the tips of  $\vec{r}(0)$  and  $\vec{r}(1)$  exactly and numerically?

- f) Write down an integral formula for the length of the curve  $\vec{r}(t)$  from  $t=0$  to  $t=1$ , and use technology to evaluate it to two decimal place accuracy. How should it compare to part e) using common sense?
- g) Evaluate the scalar tangential projection  $a_T(1)$  and the vector projections  $\vec{a}_T(1)$  and  $\vec{a}_N(1)$  of the acceleration  $\vec{a}(1) = \vec{r}''(1)$  parallel (tangential) and perpendicular (normal) to  $\vec{r}'(1)$ . How does the length of the orthogonal (normal) vector projection of  $\vec{r}''(1)$  compare to the scalar normal projection  $a_N(1) = |\vec{T}(1) \times \vec{r}''(1)|$ ?
- h) **Optional. To do only if you finish early.** Evaluate the scalar tangential and normal projections  $a_T(1)$  and  $a_N(1)$  to 3 decimal place accuracy, and the angle between  $\vec{r}'(1)$  and  $\vec{r}''(1)$  in degrees to the nearest degree. Then use these three numbers to make a rough diagram of the projection rectangle representing  $a_T(1)$ ,  $a_N(1)$  (the sides) and  $a(1) = |\vec{r}''(1)|$  (the diagonal). Label the sides with their numerical lengths, and indicate the angle too. Is the point on the curve speeding up or slowing down? Why?

► **solution**

▼ **pledge**

When you have completed the exam, please read and sign the dr bob integrity pledge and hand this test sheet stapled on top of your answer sheets as a cover page, with the first test page facing up:  
 "During this examination, all work has been my own. I have not accessed any of the class web pages or any other sites during the exam. I give my word that I have not resorted to any ethically questionable means of improving my grade or anyone else's on this examination and that I have not discussed this exam with anyone other than my instructor, nor will I until after the exam period is terminated for all participants."

Signature: \_\_\_\_\_ Date: \_\_\_\_\_