


Draw in arrows for the new basis vectors $b_1 = \langle 2, 1 \rangle$, $b_2 = \langle 1, 3 \rangle$, then extend them in each direction marking off with small bullet circles each multiple tip to tail of these along the new axes to connect with a ruler through these bullet circles to make the new coordinate axes (label them y_1, y_2 at their positive arrowhead ends). Then draw in an arrow for the vector $\langle x_1, x_2 \rangle = \langle 5, 5 \rangle$ and then draw in lines parallel to each axis from its tip to those axes to form the projection parallelogram, and read off the new coordinates $\langle y_1, y_2 \rangle$ so that $\langle x_1, x_2 \rangle = y_1 b_1 + y_2 b_2$. Next reverse the procedure for $\langle y_1, y_2 \rangle = \langle -3, 2 \rangle$ locating the point $\langle x_1, x_2 \rangle = y_1 b_1 + y_2 b_2$ drawing parallels to this point from the new axes through the tickmarks y_1 and y_2 and draw in its position vector, reading off its old coordinates. Check by matrix multiplication $x = B y, y = B^{-1} x$ that your graphical readouts are correct.



Draw in arrows for the new basis vectors $b_1 = \langle 2, 1 \rangle$, $b_2 = \langle 1, 3 \rangle$, then extend them in each direction marking off with small bullet circles each multiple tip to tail of these along the new axes to connect with a ruler through these bullet circles to make the new coordinate axes (label them y_1, y_2 at their positive arrowhead ends). Then draw in an arrow for the vector $\langle x_1, x_2 \rangle = \langle 5, 5 \rangle$ and then draw in lines parallel to each axis from its tip to those axes to form the projection parallelogram, and read off the new coordinates $\langle y_1, y_2 \rangle$ so that $\langle x_1, x_2 \rangle = y_1 b_1 + y_2 b_2$. Next reverse the procedure for $\langle y_1, y_2 \rangle = \langle -3, 2 \rangle$ locating the point $\langle x_1, x_2 \rangle = y_1 b_1 + y_2 b_2$ drawing parallels to this point from the new axes through the tickmarks y_1 and y_2 and draw in its position vector, reading off its old coordinates. Check by matrix multiplication $x = B y, y = B^{-1} x$ that your graphical readouts are correct.

$$B = \langle b_1, b_2 \rangle = \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix} \quad B^{-1} = \frac{1}{5} \begin{bmatrix} 3 & -1 \\ -1 & 2 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 5 \\ 5 \end{bmatrix}; \quad \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 3 & -1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 5 \\ 5 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 15-5 \\ -5+10 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 10 \\ 5 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix} \rightarrow \langle 5, 5 \rangle = 2b_1 + b_2$$

$$\begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} -3 \\ 2 \end{bmatrix}; \quad \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} -3 \\ 2 \end{bmatrix} = \begin{bmatrix} -6+2 \\ -3+6 \end{bmatrix} = \begin{bmatrix} -4 \\ 3 \end{bmatrix} \rightarrow -3b_1 + 2b_2 = \langle -4, 3 \rangle$$